



Assessment of Climate-Based Agricultural Water Balance in East Kalimantan Province

Akas Piningan Sujalu^{a*}

^a *Faculty of Agriculture, Univercity 17 Agustus 1945 Samarinda, Jalan Ir. H. Juanda 80 Samarinda City, East Kalimantan, 75124, Indonesia.*

Authors' contributions

The sole author designed, analyzed, interpreted and prepared the manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://prh.globalpresshub.com/review-history/1569>

Original Research Article

Received: 07/02/2024

Accepted: 11/04/2024

Published: 16/04/2024

ABSTRACT

The objectives of this study were to develop a soil water balance model for calculating the amount of water held in field storage. In dry land crop cultivation, water is the main source for plant grow and most determining limiting factor. This study conducted a Thornthwaite-Mather water balance analysis in the watershed in order to understand its hydrology capability. Components of the water balance are the existing amount of stored water, rainfall, and evapotranspiration. The result of the study show that fluctuation water balanced at East Kalimantan have 9 months water surplus with total 223 mm year⁻¹, and the dry months begins in 3 months, August tends to be the driest month in the year with total deficit value reaches 72 mm year⁻¹. Still, the annual rainfall is higher than the annual evapotranspiration.

Keywords: Water balance; rainfall; evapotranspiration; cropping.

**Corresponding author: E-mail: akaspiningansujalu@gmail.com;*

1. INTRODUCTION

“In tropical humid climates rainfall amounts are enough to satisfy dry-land agricultural production; However, this dry land agricultural schemes are very sensitive to climatic change and the productivity is linked to precipitation patterns. Soil properties in the tropics become very important in their function as a buffer system for the development of food security systems, especially regarding the presence of water, with the irregularity of rainfall. In dryland crop cultivation, water is the most determining limiting factor and rain is the main source for plant growth. The variation in rainfall, both in amount, intensity and time may hamper the prediction of planting time or planting pattern management caused by fluctuated water availability” [1,2,3].

“This water fluctuation often occurs in the dryland, so that efficient and effective land planning is needed through soil water balance information. This soil water balance is a water balance for general agricultural land use” [4]. “This balance sheet is useful in considering the suitability of agricultural land, arranging planting and harvesting schedules as well as regulating irrigation water provision in the right amount and time. Soil water balance is a balance between groundwater stored as a reserve in the soil, derived from irrigation water and rainfall, with potential water loss in the form of drainage, surface runoff, evaporation and transpiration” [5].

“Actual water use in annual water balances is changing over time as cropping patterns and cropping intensities shift and change, sometimes in response to available water resources. This water fluctuation often occurs in the dryland, so that efficient and effective land planning is needed through soil water balance information. Agriculture accounts for about 70% of the freshwater withdrawals in the world, while consumptive use of water in agriculture (water that is evaporated on irrigated fields) accounts for about 90% of all of the water that is evaporated as a result of human intervention” [6, 7,8].

“So important is the availability of water for procurement planning and improvement of agricultural support facilities, that the success of a farm management is also determined by how far the availability of water can be distributed by physical facilities” [1,8,9,10]. For this reason, research was conducted to analyze land water balance based on normal rainfall in East

Kalimantan Province. Each component of the water balance and at a certain time or period desired according to its needs. This water balance has an accumulative nature, while the unit of each component of water balance is the height or range of water (mm or cm) and the unit of time is daily, weekly, monthly and yearly [10,11,12,13]. The study is conducted by analyzing the availability of soil water in the form of graphs presented monthly, from January to December on a climatological base.

2. MATERIALS AND METHODS

2.1 Research Location

The Province of East Kalimantan, Indonesia with the capital city of Samarinda before the division of the region has an area of East Kalimantan province possesses an area of 16,732,065.18 hectare, with land area of 12,734,691.75 hectare. Located between 113°44' East Longitude and 119°00' East Longitude and between 2°33' North Latitude and 2°25' South Latitude. The land area of East Kalimantan Province has an undulating topography with a slope that is from steep to highest, with land elevations ranging from 0–1,000 m above sea level.

East Kalimantan tropical climate and has two seasons, dry and rainy seasons. The dry season usually occurs in May to October, while the rainy season in November to April. This situation continued every year interspersed with transitional season in certain months. Moreover, because of its location on the equator, the climate in East Kalimantan are also affected by wind monsoon, monsoon wind is November–April west and east monsoon winds from May to October. In recent years, the situation in East Kalimantan season is sometimes erratic. In the months that it is supposed to rain, there is no rain at all, or vice versa in the months that should be dry it rains for a much longer time in general, East Kalimantan hot climates with temperatures ranged from 21.6 °C in October to 35.6 °C in September. A side from being a tropical area with extensive forests, the average humidity between 83% and 87%. The lowest air humidity observed by the meteorological station happens in a few months with 82% humidity, while the highest occurred in February with 91% humidity.

In this study using historical data on rainfall the research object observed was the condition and extent of forest cover along with the condition of




Fig. 1. The East Kalimantan Province study area is located on Borneo

climate elements "in situ" from 2000 to 2020 from 76 Meteorological Stations in East Kalimantan Province, data was obtained from Agriculture and Horticulture Department, Meteorological Climatological and Geophysical Agency (BMKG) as the main source of weather data. The method used in this research is a quantitative descriptive method with comparative analysis techniques of exploration of the area data and the characteristics of the climate elements that are owned.

2.2 Analysis of Land Water Balance

Water balance analysis is expressed in the form of integral equations by simplifying some equations, so that the water balance of an agricultural project area can be expressed in the form of equations:

$$P = \text{ETA} \pm \Delta \text{SWC} \pm \text{Ro}$$

P = Rainfall (mm)
 ETA = Actual Evapotranspiration
 Δ SWC = Soil Water Content
 Ro = Run off

The water balance affects how much water is stored in a system. The general water balance in the UK shows seasonal patterns. In wet seasons precipitation is greater than evapotranspiration which creates a water surplus. Ground stores fill with water which results in increased surface runoff, higher discharge and higher "water table"

levels. Precipitation provides part of the water crops need to satisfy their transpiration requirements. The soil stores part of the precipitation water, which is later evaporated or transpired by plants [8,9,10,11,13,14]

Thornthwaite-Mather water balance equation uses the soil moisture capacity to estimate water budgets [12]. The parameters needed for using this method include:

1. difference between precipitation and potential evapotranspiration (P-PE)
2. accumulated potential water loss (APWL)
3. available water capacity (AWC)
1. values during periods when P-PE is negative. This running represents the total amount of
4. actual evapotranspiration (AE)
5. deficit and surplus of the water budget 7 runoff estimation

"Accumulated Potential Water Loss (APWL) is a condition in which the number of potential evapotranspiration is greater than the amount of rainfall, which indicates a dry month. The equation for Accumulated Potential Water Loss (APWL) is as follows: $APWL = P - PE$, sum of the daily P - PE values during periods when P - PE is negative. This running sum represents the total amount of unsatisfied potential evapotranspiration to which the soil has been subjected" [11,12,13,14].

3. RESULT AND DISCUSSION

3.1 Climate Status in East Kalimantan

In this study, climatological data (including irradiation duration, irradiation intensity, monthly average air temperature and monthly average air humidity) obtained from 76 weather observing stations were collected, as follows:

3.2 Land Water Balance

Monthly water balance analysis of the above-mentioned results shows that these areas have surpluses during the eight Monthly water balance analysis of the above-mentioned results shows that these areas have surpluses during the eight months that occur in the month periods from January to June, and October - December.

In addition to experiencing a monthly water surplus, the eastern Kalimantan Province also experienced a monthly cumulative water deficit in the 3-month period from July to September. APWL values are utilised for Actual Evapotranspiration (ETA) and the absence of rainwater supply causes the deficit (Fig. 2). The effect of this water deficit depends on the water-holding capacity of the soil, which differs for each soil type influenced by clay content (texture), organic matter and topography [13,15]. That

period was the dry season characterised by a water deficit in the East Kalimantan province, and the dry season occurred evenly across the region in July, and the impact of the drought amplified until it reached its peak in August to September. The lack of rainfall during July to August was followed by relatively high air temperatures leading to a water deficit in the active root zone causing a disruption of the soil water balance. Despite this, agricultural lands in the eastern Kalimantan province theoretically do not yet require additional irrigation water as the water status in the soil is still >50% Available Soil Water Content [10,13,14,16].

The results of the analysis of monthly land water balance of the East Kalimantan Province region were subjected to evapotranspiration potential (ETP) analysis which was calculated based on monthly average air temperature data. It is known that this area has a surplus for 9 months which occurs in the period January - June and in October - December the total water surplus reaches 223 mm / year. In addition to having a water surplus, the area normally experiences a monthly cumulative water deficit due to rainfall received being lower than the corrected evapotranspiration rate (CH < corrected ETP). Based on the data analysis in the appendix Table A 1, the water deficit occurred in the July - September period which totalled 72 mm/year.

Table 1. Monthly Average Climate Element Data for East Kalimantan Province (113°44'E and 119°00'E, and between 2°33'N and 2°25'S)

Climate parameter	Months											
	1	2	3	4	5	6	7	8	9	10	11.	12.
Rainfall (mm/bulan)	166	178	166	184	165	145	126	118	107	155	130	205
Rain days (days)	12	11	12	14	12	11	9	10	9	11	11	12
Temperature (°C)	26.8	26.7	26.6	26.4	26.4	26.3	26.2	26.0	26.0	25.8	25.7	26.2




Fig. 2. Water balance East Kalimantan province

Thus resulting in potential groundwater depletion. Under such conditions of circulating water availability and depletion, water needs for farming areas must be added through physical means of irrigation networks [16,18].

When viewed from the aspect of water requirements for the development of physical facilities, the rainfall on dry land as illustrated in the Monthly Land Water Balance (Cumulative) in addition to providing a variety of advantages, including very possible to develop the fisheries sector, especially inland fisheries (freshwater ponds) but also shows the existence of constraints - constraints that are difficult to overcome manually [7,8,9]. The water surplus experienced during the 9 months can cause a high intensity of disturbance to the construction of physical facilities, especially disturbance from high levels of erosion and heavy water discharge that must be removed from the farming area. Because surplus water that cannot be released smoothly will cause various things that are detrimental to farming [17] Among them are the high chances of the emergence of disturbances in percaran due to decay, especially in land management for the purpose of tubers [18,19].

Some important information that can be taken for technical purposes operations in the field is that in months where there is a consecutive water surplus, it is necessary to make a drainage/disposal channels to drain excess water, especially if excess water, especially if soil conditions do not conditions are not favorable for rapid infiltration [9, 14]. As for the deficit period (July-September) are still possible to do business crop cultivation because groundwater conditions are still favorable for the business and remain in an available state. In September and October it is recommended to be careful in planting activities especially planting short-rooted crops. It is possible to carry out planting activities planting activities then during these months must be supplied with water. Watering (irrigation) measures are still needed so that plants can grow and develop properly according to their genetic potential.

In addition, considering the topographical conditions of the region, it is possible to build several dams or reservoirs in East Kalimantan Province, which have various functions. With the main function being to accommodate the considerable water surplus in the region, as well as a water reserve during water deficit months

that can be utilised for various purposes including drinking water. This is very possible if the irrigation channels that function to dispose of excess water are not well organised and maintained.

4. CONCLUSION

Based on the results of the Land Water Balance analysis and the overall description in the discussion as previously presented, it can be concluded that the province of East Kalimantan experiences rainfall every month, even in dry years. Rainfall in a normal climatic year ranges from 107 mm/month - 205 mm/month with 9-12 rainy days. The region has a water surplus experiencing 9 months of groundwater surplus and 3 months of groundwater deficit resulting in Accumulation Potential water loss although the groundwater status is still above field capacity.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Azmeri A, Yulianur A, Zahrati U, Faudli I. Effects of irrigation performance on water balance: Krueng baro irrigation scheme (Aceh-Indonesia) as a case study. *Journal Water Land Deviation*. 2019;42:12-20.
2. Pathak P, Sudi R, Wani SP, Sahrawa KL. Hydrological behavior of Alfisols and Vertisols in the semi-arid zone: Implications for soil and water management. *Agricultural Water Management*. 2015;118:12-21
3. Reichardt K, Timm LC. The water balance in agricultural and natural Systems. *Soil, Plant and Atmosphere*. 2020:289-312.
4. Maihemuti B, Simayi Z, Alifujiang Y, Aishan T, Abliz A, Aierken G. Development and Evaluation of the soil water balance model in an Inland Arid Delta Oasis: Implications for sustainable groundwater resource management. *Global Ecology and Conservation*. 2021;25:e01408.
5. Pereira LS, Paredes P, Jovanovic N. Soil water balance models for determining crop water and irrigation requirements and irrigation scheduling focusing on the FAO56 Method and the dual Kc approach. *Agricultural Water Management*. 2020;241:106357

6. FAO. The State of Food and Agriculture 2013. Food and agriculture organization of the United Nations.
7. Babaei M, Mahmoodzadeh D, Ketabchi H, Saadi T. Estimation of water balance components and analysis of variations using modeling and remote Sensing Approaches (Hashtgerd Study Area, Alborz Province). *Iran-Water Resources Research*. 2022;18(1):31–53.
8. do Vale TMC, Spyrides MHC, Cabra JB, IJ, Andrade LMB, Bezerra BG, Rodrigues DT, Mutti PR. Climate And Water Balance Influence On Agricultural Productivity Over The Northeast Brazil. *Theoretical and Applied Climatology*. 2023; 155:879–900
9. Koivusalo H, Turunen M, Salo H, Haahti K, Nousiainen R, Warsta L. Analysis of water balance and runoff generation in high Latitude Agricultural fields during mild and cold winters. *Hydrology Research*. 2017;48 (4):957–968.
10. Rusli SR, Weerts AH, Taufiq A, Bense VF. Estimating Water Balance Components and their uncertainty bounds in highly Groundwater-Dependent and Data-Scarce Area: An example for the Upper Citarum basin. *Journal of Hydrology. Regional Studies*. 2021;37:100911
11. Chen J, Chen S, Ma F, Chen J. Analysis of water balance of hulun lake based on digital remote sensing images. *Water Resources Protection*. 2020;36;73–79.
12. Nugroho AR, Ichiro Tamagawa, Almaika Riandraswari, and Titin Febrianti. Thornthwaite-Mather water balance analysis in Tambakbayan waterhed, Yogyakarta, Indonesia. 2019. *MATEC Web of Conferences* 280;05007
13. Lim CS, Chae HS. A Study on Association. 2007;40:555–570.
14. Mammoliti E, Fronzi D, Mancini, A, Valigi D, Tazioli A. Waterbalance, a WebApp for Thornthwaite-Mather Water Balance Computation: Comparison of applications in two European Watersheds. *Hydrology*. 2021;8(1);34-44
15. Chaogui Lei, Paul D. Wagner, Nicola Fohrer. Effects of land cover, topography, and soil on stream water quality at Multiple Spatial and seasonal scales in a German Lowland Catchment. *Ecological Indicators*. 2021; 120:106940
16. Nath TN. Soil Texture and total organic matter content and its influences on soil water holding capacity of some selected Tea Growing Soils in Sivasagar District Of Assam, India. 2014. *International Journal of Chemical Sciences* 12(4):1419-1429
17. O'Geen AT. Soil Water Dynamics. *Nature Education Knowledge*. 2013;4(5):9
18. Turunen M.W.L. Paasonen-Kivekäs., M. Nurminen J. Mylly M. Alakukku L. Äijö H. Puustinen and M. Koivusalo H. Modeling Water Balance And Effects Of Different Subsurface Drainage Methods On Water Outflow Components In A Clayey Agricultural Field In Boreal Conditions. *Agricultural Water Management*. 2013;121: 135–148
19. Yang HK. Water balance change of watershed by climate change. *Journal of Korea Geografic Society*. 2007;42;405–420.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://prh.globalpresshub.com/review-history/1569>